

IMPORTANT INFORMATION

Warning signs will be placed at all main access points to the operational area and everyone must follow the cautions on the signs. There's no health risk when using this area as long as you follow these instructions:

Do not handle any bait or allow children to wander unsupervised. Cereal baits containing 1080 are dyed green.

Do not hunt or take game from within a two kilometre radius of the operational area for human or pet consumption. It's an offence

to sell meat products that have been exposed to 1080. Hunting can resume approximately four months following the control work, or after two months if 100mm of rain has fallen.

Please observe these rules whenever you see warning signs about the pesticide. Warning signs indicate that pesticide residues may still be present in baits or animals. When the signs are officially removed, you can resume normal activities in the area. Free dog muzzles will be provided on request. Please contact OSPRI on 06 3532710 to obtain a muzzle.



Do not bring dogs into the area until the warning signs have been officially removed.

Dogs are particularly susceptible to 1080. They must not be allowed access to bait or poisoned carcasses which remain toxic to dogs until they have fully decomposed.



WHAT TO DO IF YOU SUSPECT POISONING

Contact your local hospital or doctor, or dial 111

National Poisons Centre
0800 POISON (764 766)

If a domestic animal is poisoned, contact a local veterinarian.



FURTHER INFORMATION

OSPRI
North Island office,
Palmerston North
P 06 353 2710
E vector.SNI@ospri.co.nz
tbfree.co.nz | ospri.co.nz

Contractor
BioWorks
Masterton
P 06 378 2484
E Graeme.Butcher@gw.govt.nz

For more information on controlling bovine TB and why 1080 is used in New Zealand, visit 1080facts.co.nz



SOUTHERN RIMUTAKA AERIAL BOVINE TB CONTROL OPERATION

OSPRI'S TBfree PROGRAMME


The TBfree programme aims to manage and eventually eradicate bovine tuberculosis (TB) from New Zealand's farmed cattle and deer and wild animal populations. We use possum control, along with regular herd testing and movement restrictions, to achieve this objective. Information gathered from wild animal surveys, recent and historic findings of TB in wild animals, herd testing results and the operational history of the region is used when planning operations.



TARGETING POSSUMS

To eradicate bovine TB, possum numbers need to be kept extremely low – around one to two animals every 10 hectares. Monitoring has shown that control work is needed in the Southern Rimutaka area to reduce the possum population and minimise the risk of the disease spreading through wild animal populations and onto farmed cattle and deer. Possum control has dramatically reduced the number of infected herds in the Greater Wellington region. Controlling disease prevents livestock production losses and protects the world-leading reputation of New Zealand's dairy,

HOW WE CONTROL TB



beef and deer products. This area has not been treated by OSPRI's TBfree programme before.

WHAT TO EXPECT

The operation will begin with the distribution of non-toxic, tan-coloured cereal pellets by helicopter. This "pre-

feed" gives possums a taste for the pellets and overcomes bait shyness. One to two weeks later (weather permitting), toxic, green cereal pellets coated with deer repellent – each containing 0.15% biodegradable sodium fluoroacetate (also known as 1080) – will be applied by helicopter at a rate of two kilograms per hectare.



TBfree is an OSPRI programme



tbfree.co.nz



0800 482 463



TBfree is an OSPRI programme



tbfree.co.nz



0800 482 463



That’s about one bait to every 60 square metres. The Southern Rimutaka operation will be subject to strict safety, quality-assurance and monitoring requirements.

Advanced GPS navigational equipment will be used to ensure the pellets are accurately placed and identified exclusion zones avoided.

THE METHOD

A vast majority of possum control in the region is done by local contractors using ground-based traps and hand-laid toxins. The remaining area is controlled using aerially applied pellets containing biodegradable 1080. Aerial control is highly-efficient, cost-effective and has historically been extremely successful at knocking possum numbers down to very low levels. It is preferred in areas like Southern Rimutaka due to the rugged nature of the terrain. The Parliamentary Commissioner for the Environment also supports aerial



Baits covered in deer repellent. The non-toxic pre-feed pellets are brownish-tan while the toxic baits are green.

control. The commissioner completed an extensive review into the use of 1080, in which she strongly endorsed its continued use in New Zealand.

Please visit pce.parliament.nz to read this report.

WHAT HAPPENS NOW

All landowners/occupiers within and adjacent to the proposed control area will be contacted and visited by TBfree contractors ahead of the operation. They will discuss boundary issues, water supply safety and the management of any risks to dogs and livestock. Consents from Greater Wellington Regional Council, the Department of Conservation and Ministry of Health are required for this operation. Affected landowners/occupiers will be contacted again before the operation starts, notices will be published in local newspapers and warning signs will be placed at all likely access points to the operational area.

BIODIVERSITY BENEFITS

The operation will have additional conservation benefits for native birds and bush. Possums eat the forest canopy and prey on native birdlife, including eggs and chicks. Biodegradable 1080 is also extremely effective at controlling other introduced predators such as ship rats and stoats.



Boundaries are indicative only and will change after consultation